



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

377. Proposed by W. D. CAIRNS, Oberlin College.

It is required to find a curve of the form $y = x(x - a)(x - b)$ such that the abscissas of the maximum and minimum values, as well as a and b , shall be positive integers.

MECHANICS.

When this issue was made up, no solutions had been received for numbers 289, 292-3, 295-299, 301, and 303.

302. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

A ball is projected from a given point at a given inclination β towards a vertical wall; determine the velocity so that after striking the wall the ball may return to the point of projection.

NUMBER THEORY.

When this issue was made up, no solutions had been received for numbers 215-16, 218, 220, and 223-226.

226. Proposed by ELBERT H. CLARKE, Purdue University.

If $0!$ is taken equal to 1, and if k is any positive integer greater than or equal to 2, show that

$$\sum_{n=0}^{\infty} \frac{n!}{(k+n)!} = \frac{1}{(k-1)!} \cdot \frac{1}{(k-1)!}.$$

227. Proposed by R. P. BAKER, University of Iowa.

Show that every rational number can be expressed as a finite sum $\sum_{n=m}^{n=m+k} \frac{a_n}{n}$, where a_n is either 0 or 1 and m is any positive integer.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

409. Proposed by C. E. GITHENS, Wheeling, W. Va.

Find integral values for the edges of a rectangular parallelepiped so that its diagonal shall be rational.

II. SOLUTION BY ARTEMAS MARTIN, Washington, D. C.

On pp. 269-273 of the October, 1914, MONTHLY, W. C. Eells has solved a different problem from the one proposed. In making $x^2 + y^2 = \square$, he adds another condition not required.

Let x, y, z be the edges and d the diagonal of the parallelepiped; then we have to satisfy the equation

$$x^2 + y^2 + z^2 = d^2.$$

It is not necessary that $x^2 + y^2$ be a square. Let us assume $x = a$, $y = b$, $z + c = d$, and we have

$$a^2 + b^2 + z^2 = (z + c)^2 = z^2 + 2cz + c^2,$$

which immediately gives

$$z = \frac{a^2 + b^2 - c^2}{2c} \quad \text{and} \quad d = \frac{a^2 + b^2 + c^2}{2c}.$$